

Current Journal of Applied Science and Technology

Volume 44, Issue 1, Page 87-101, 2025; Article no.CJAST.129848 ISSN: 2457-1024 (Past name: British Journal of Applied Science & Technology, Past ISSN: 2231-0843, NLM ID: 101664541)

Metritis in Dairy Cattle: Diagnostic Challenges, Economic Impacts, and Emerging Insights

Gayathri S. L. ^{a*}, Bhakat M. ^b and Mohanty T. K. ^a

^a Livestock Production Management Division, ICAR-National Dairy Research Institute, Karnal-132001, Haryana, India.

^b Animal Physiology and Reproduction Division, ICAR-Central Institute for Research on Goats, Makhdoom, Farah-281122, Matura, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/cjast/2025/v44i14478

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/129848

Review Article

Received: 14/11/2024 Accepted: 16/01/2025 Published: 18/01/2025

ABSTRACT

Metritis, an inflammatory uterine disorder occurring within 21 days postpartum, poses significant global challenges to dairy herd management. This review comprehensively examines the incidence, diagnostic criteria, clinical manifestations, economic implications, and emerging management strategies for metritis in dairy animals. The classification of metritis, distinguishing between puerperal metritis and delayed uterine involution, is discussed alongside diagnostic approaches such as vaginal discharge scoring, cytology, and rectal temperature evaluation. Metritis incidence varies widely, ranging from 10% to 36%, with early postpartum diagnosis being critical to minimizing its impact. The economic burden of metritis is profound, with affected cows experiencing reduced milk yield, impaired reproductive performance, extended calving intervals, and increased culling

Cite as: S. L., Gayathri, Bhakat M., and Mohanty T. K. 2025. "Metritis in Dairy Cattle: Diagnostic Challenges, Economic Impacts, and Emerging Insights". Current Journal of Applied Science and Technology 44 (1):87-101. https://doi.org/10.9734/cjast/2025/v44i14478.

^{*}Corresponding author: E-mail: gayathrisherlylal@gmail.com;

rates. Despite its clinical significance, diagnosing metritis remains challenging due to inconsistent presentation of fever and subjective assessment of clinical signs. Non-invasive diagnostic techniques, such as infrared thermography (IRT), are highlighted as promising tools for early detection, offering a stress-free and practical approach to monitoring rectal temperature and systemic health. Recent advances in molecular research have uncovered genetic variants and regulatory pathways, such as cholesterol metabolism and the PI3k-Akt signalling pathway, associated with disease susceptibility and progression. These insights provide potential therapeutic targets and opportunities for precision management strategies. The present review accentuates the critical need for standardized diagnostic criteria, innovative non-invasive technologies, and integrative management approaches to address metritis effectively. Continued research and the application of cutting-edge technologies are essential to mitigate the economic and welfare impacts of metritis, ultimately improving herd health, productivity, and sustainability in the dairy industry.

Keywords: Metritis; dairy animals; diagnosis; temperature.

1. INTRODUCTION

Metritis is a significant postpartum uterine disorder in dairy cows, typically occurring within 10 to 21 days after calving, characterized by uterine inflammation often accompanied by systemic illness, with or without fever. The condition exerts considerable negative effects on dairy production systems globally, impairing milk yield, reproductive performance, and cow welfare while increasing the risk of early culling (Ribeiro et al., 2013; Stojkov et al., 2015; Dubuc et al., 2011). The economic burden associated with metritis is substantial, stemming from reduced milk production-up to 5 kg/day-prolonged calving-to-conception intervals, increased treatment costs, and higher culling rates (Giuliodori et al., 2013; Daetz et al., 2016; Barragan et al., 2018). Additionally, metritis has been closely linked to poor animal welfare outcomes, emphasizing the need for early and effective management strategies (Neave et al., 2018).

Despite its global prevalence, ranging between 10% and 36% of dairy cows (Sheldon et al., 2006; De Oliveira et al., 2020), metritis diagnosis remains inconsistent. Traditional diagnostic criteria often categorize puerperal metritis based on an abnormally enlarged uterus, fetid uterine discharge, and systemic signs like However, variations in disease fever. presentation complicate this approach, as many cows with fetid discharge do not exhibit fever, and some cases involve delayed uterine involution without systemic symptoms (Sheldon et al., 2006; Martinez et al., 2012). Garzon et al. (2022) emphasise in their study for an urgent need for a unified framework and robust diagnostic protocols to address these discrepancies.

Conventional diagnostic methods, such as discharge scoring and cytological vaginal evaluations, have been widely used but are limited by subjective interpretation and laborintensive procedures (Jeon et al., 2016). Advanced laboratory techniques. including microbiology and polymerase chain reaction (PCR), offer precision but are time-consuming, expensive, and reliant on specialized facilities, restricting their utility in field conditions (Garzon et al., 2022). In this context, the development of portable. non-invasive diagnostic tools represents a paradigm shift in metritis detection and management.

Emerging technologies like the Metrisor device (Risvanli et al., 2024) and infrared thermography (IRT) (Macmillan et al., 2019) exemplify this shift. The Metrisor utilizes gas sensors to detect intrauterine gases, providing a rapid, accurate, and portable diagnostic solution tailored for field use. Similarly, IRT offers a non-invasive method to identify early postpartum illnesses through surface temperature variations, enhancing disease detection without the need for invasive procedures. These tools hold promise for addressing the limitations of current methods, enabling real-time, scalable diagnostics directly on farms.

This review aims to bridge the existing gaps in metritis detection and management by critically definitions, classifications, evaluating its prevalence, and economic impacts. In doing so, synthesizes insights from emerging it technologies and recent research to propose novel, field-adaptable diagnostic frameworks, By practical emphasizing the feasibility and transformative potential of these advancements, this review seeks to support the development of comprehensive strategies to improve herd

health, productivity, and animal welfare while mitigating the economic burden of metritis on the dairy industry.

2. METRITIS- DEFINITION AND CLASSIFICATION

Metritis is a significant postpartum uterine disorder characterized by an abnormally enlarged uterus and the presence of fetid, waterv. reddish-brown uterine discharge, typically occurring within the first 21 days after parturition (Sheldon et al., 2006; Stojkov et al., 2015). Its prevalence peaks within the first ten days postpartum, making it a critical concern for dairy herd management. Depending on clinical presentation and severity, metritis is classified into clinical metritis and puerperal metritis (PM), with severity ranging from subclinical to lifethreatening toxemic conditions.

Presentation Challenges Clinical and in Definition: The characteristic features of metritis include an enlarged, flaccid uterus and discharge varying from watery red-brown to viscous purulent with a distinct fetid odour (Sheldon et al., 2006). However, a unified definition of metritis remains elusive due to variations in clinical signs and diagnostic criteria among studies. PM, a severe form of metritis, involves inflammation of all uterine layers, systemic illness, and fetid discharge. Systemic symptoms often include pyrexia (>39.5°C), inappetence, dullness, and decreased milk production, typically manifesting between 7 to 14 days postpartum (Olson et al., 1986; Sheldon et al., 2006, 2009).

The lack of standardization in diagnostic criteria poses challenges for clinical and research applications. For instance, fever thresholds for PM diagnosis vary across studies, ranging from 39.2°C to 39.7°C (Drillich et al., 2001; Sheldon et al., 2006). Moreover, disparities in describing vaginal discharge—using terms such as redbrown, chocolate, or yellow for color and fetid or putrid for odor—highlight the subjective nature of

these evaluations (Garzon et al., 2022). Such inconsistencies can lead to discrepancies in diagnosis, treatment recommendations, and antimicrobial use, underscoring the need for a standardized framework.

Microbial Etiology and Disease Complexity: Metritis is a multifactorial disease predominantly caused by polymicrobial infections. Kev pathogens implicated in postpartum uterine infections include Escherichia coli. Trueperella pyogenes, Fusobacterium necrophorum, and Bacteroides which often interact spp., synergistically with other bacteria such as Helcococcus, Filifactor. Porphyromonas, Peptoniphilus, Peptostreptococcus, Campylobacter, and Prevotella to exacerbate disease severity (Jeon et al., 2015). Also, Cunha et al. (2018) reported that metagenomic studies question the traditional focus on Escherichia coli and *Trueperella pyogenes* as primary pathogens in metritis. Instead, they highlight the importance Fusobacterium necrophorum. Prevotella of melaninogenica, and emerging pathogens such as Bacteroides pyogenes, Porphyromonas levii, and Helcococcus ovis in the disease's etiology. The disease's progression depends on factors such as pathogen virulence, uterine environment, and immune response, making it a complex and multifaceted condition.

Puerperal Metritis and Disease Complex: Puerperal metritis (PM), also referred to as acute puerperal metritis (APM), occurs within ten days postpartum and is characterized by systemic illness, including fever ≥39.5°C, fetid uterine discharge, and an enlarged uterus (Sheldon et al., 2006; Machado et al., 2014). The systemic manifestations of PM, such as toxaemia, depression, and reduced productivity, make it a significant health concern. Toxic puerperal metritis (TPM), a severe subtype of PM, presents with increased rectal temperature, fetid vulvar discharge. and uterine atony, requiring immediate intervention to prevent septicemia and mortality (Smith et al., 1998).



Fig. 1. Classification of metritis

2.1 Fever in Metritis: Mechanisms, Implications, and Immunological Responses

The treatment protocol for metritis remains consistent, irrespective of the presence or absence of pyrexia (McLaughlin et al., 2012). However, metritis-associated fever introduces unique physiological and clinical implications. Both febrile and afebrile metritic cows exhibit reductions in milk production and an increased likelihood of developing clinical endometritis (Benzaguen et al., 2007). However, a significant decline in reproductive performance is observed primarily in metritic cows with fever, potentially due to the systemic inflammatory burden (Giuliodori et al., 2013). Recovery rates for febrile metritic cows are notably lower compared to their afebrile counterparts, which may be attributed to variations in uterine microbiota that influence the response to antibiotic therapy (Lima et al., 2014).

Fever in metritic cows arises from the activation of innate immune pathways triggered by uterine bacterial infections. The inflammatory response is mediated by the induction and release of endogenous cytokines such as tumour necrosis factor-alpha (TNF α), interleukin-1 beta (IL-1 β), and interleukin-6 (IL-6), along with chemokines like RANTES and MCP-1. These mediators activate polymorphonuclear and mononuclear leukocytes, culminating in the synthesis of prostaglandin E (PGE), which triggers fever via the central nervous system (Netea et al., 2000). Among bacterial components, lipopolysaccharides (LPS) from Gram-negative bacteria serve as the most potent fever-inducing agents, though peptidoglycans from Grampositive bacteria can also elicit febrile responses (Tavares et al., 2005).

In metritis, fever is commonly associated with uterine infections dominated by Gram-negative bacteria such as *Fusobacterium* spp. and *Bacteroidetes* spp., known for their ability to stimulate robust inflammatory responses (Jeon et al., 2015). Similarly, neutrophils play a pivotal role in propagating inflammation and clearing bacterial infections during uterine infections (Sheldon et al., 2009). However, compromised neutrophil function has been observed in dairy cows during the periparturient period, extending from one week before to three weeks after calving, which predisposes cows to metritis (Hammon et al., 2015).

Immunosuppression during the transition period creates a window of vulnerability, allowing bacterial infections to establish. However, not all cows exposed to pathogenic bacteria develop metritis, suggesting that bacterial load and virulence are primary determinants of disease severity and febrile response (Jeon et al., 2015, 2016). These findings underscore the complexity of metritis pathogenesis and highlight the need for targeted therapeutic strategies to mitigate the systemic effects of febrile metritis and improve recovery outcomes.

2.2 Incidence and Prevalence of Metritis

Puerperal metritis continues to be a major health concern in postpartum dairy cows, with prevalence rates ranging from 5% to 20% within herds worldwide (LeBlanc, 2014) and metritis in the range of 8 % to 40% (Silva et al., 2024). Despite a global focus on the condition, there is still a lack of extensive research in some regions, such as India. Benzaguen et al. (2007) conducted a study on a commercial dairy farm with 1,000 lactating cows in northeast Florida, finding a 21% incidence of puerperal metritis. The study also revealed seasonal effects, with primiparous cows calving during the warm season having lower odds of developing metritis compared to those calving during the colder months. A more recent study by Torres et al. (2020) in northern Mexico found a 16% prevalence of metritis among 2,500 Holstein cows, with 11.3% of cows showing clinical endometritis following retained fetal membranes (RFM).

Globally, the prevalence of metritis varies across different regions. For example, Kasimanickam et al. (2004) reported a high incidence of 45% in the United States (Kasimanickam et al., 2004), while studies from Canada (13.5%) (Dubuc et al., 2010), Argentina (38%) (Plontzke et al., 2010), New Zealand (29%) (McDougall et al., 2011) and Belgium (27.8%) (Bogado et al., 2016) also show significant variation. In Argentina, Giuliodori et al. (2013) found a 29.7% incidence of puerperal metritis and a 9.6% incidence of clinical metritis in a study of 303 Holstein-Friesian cows. The study highlighted those multiparous cows had lower odds of developing metritis compared to primiparous cows. Factors such as abnormal calving, high pre-partum non-esterified fatty acids (NEFA) levels, and lower pre-partum insulin-like growth factor-1 (IGF-1) were associated with increased risk. Zwald et al. (2004) documented a 21% incidence of metritis, including RFM, from a large dataset involving 97,318 cow records in the United States. Gaddis et al. (2012) reported that, based on health records from 1996 to 2009 in the United States, the incidence of metritis in cows varied between 2% and 36%, with an average prevalence of 13%.

More recent studies have focused on identifying key risk factors. Kaniyamattam et al. (2020) observed a consistent 28% incidence of metritis across both first and subsequent lactations in US dairies. De Vries (2017) proposed that retaining a greater proportion of calves could help reduce metritis incidence by increasing the annual cull rate. McCarthy and Overton (2018) found that mild metritis occurred in 24.7% of cases, with severe metritis found in 3.5%, in a study involving a Holstein herd with a one-year calving interval. Misclassification bias is believed to underestimate the true association between metritis and its economic impact, as well as its effects on milk production (Gaddis et al., 2012; McArt et al., 2015; Liang et al., 2017). Newer research continues to refine our understanding of the complex interactions between environmental, physiological, management and factors influencing metritis prevalence. The need for more standardized diagnostic criteria across regions is critical in improving the accuracy of prevalence reports and in mitigating the disease's economic impact on the dairy industry.

2.3 Production and Economic Loss Due to Metritis

Metritis is a major cause of economic loss in dairy farming due to various factors such as treatment costs, milk withdrawal leading to lost milk production, reduced reproductive performance, and increased premature culling (Heikkilä et al., 2012; Lima et al., 2019). Several variables, including the milk yield, milk price, and the duration of milk withdrawal during treatment influence the economic burden of metritis. Puerperal metritis, or early metritis, significantly decreases milk yield and feed intake (Huzzey et al., 2007), while also negatively impacting reproductive efficiency (Melendez et al., 2004). The average cost per metritis case has been reported at approximately \$350 (Overton and Fetrow, 2008), with more recent estimates suggesting costs of \$511 per case in dairy herds maintained in the USA (Pérez-Báez et al., 2021). This increase highlights the escalating financial burden of metritis and the need for more effective management strategies to mitigate these costs. Beyond direct treatment costs, metritis affects

the dairy industry by reducing milk production, impairing reproductive performance, and increasing culling rates, further compounding the financial losses (Lima et al., 2019; Machado et al., 2020). Despite variations in estimations across studies, the economic toll of metritis is clear, emphasizing the importance of continued research to evaluate its impact across diverse dairy farming systems (Mahnani et al., 2015).

Metritis also incurs additional costs related to treatment, which typically range from \$46 to \$101 per case, depending on factors like cow parity and the choice of antibiotic (Lima et al., 2019). Previous studies reported varying cost estimates, with Bartlett et al. (1986) estimating a total cost per case of \$106. More recent studies have found cost ranges between \$329 and \$386 per case (Overton and Fetrow, 2008). These discrepancies reflect different treatment approaches, disease management practices, and farm conditions, suggesting the need for more comprehensive studies that consider these factors across various farm settings to accurately quantify the economic burden of metritis in the dairy industry. The detrimental effects of metritis on milk production and fertility are well documented. Cows suffering from metritis generally experience decreased milk production and poorer fertility, and they are more likely to be culled (Stangaferro et al., 2016). Lima et al. (2019) estimated that economic losses per metritis case ranged from \$267 to \$410 due to the combined effects of antibiotic treatment and the adverse impacts on milk production, reproduction, and culling. Drillich et al. (2001) similarly calculated a loss of \$380 per affected COW.

Machado et al. (2020) assessed how cow-related factors influence the cure rate and associated economic outcomes of metritis. They found that cows with higher levels of haptoglobin (Hp >0.54 mg/mL), which is a marker of inflammation, produced less milk in the early stages of lactation. Specifically, metritic cows that were not treated with ceftiofur had significantly reduced milk production compared to non-metritic cows. Additionally, metritis negatively affects reproductive efficiency, increasing the number of davs between calving and first service (Fourichon et al., 2000), and requiring more resources to induce pregnancy.

Culling decisions on dairy farms are influenced by an animal's reproductive health, with metritis increasing the risk of premature culling, particularly among multiparous cows (Gröhn et al., 2003). Wittrock et al. (2011) found that multiparous cows with metritis ate less during the first three weeks after calving, resulting in reduced milk production and an increased likelihood of culling by 305 days. This long-term reduction in milk production may be partly attributed to a decline in feed intake, which limits the energy available for milk synthesis (Bell and Roberts, 2007; Huzzev et al., 2007). Interestingly, primiparous cows with metritis were found to have a faster immune response to infection and recovered more quickly than multiparous cows (Waldron et al., 2006). Giuliodori et al. (2013) reported that cows with metritis (both clinical and puerperal) produced less milk than healthy cows during early lactation but more milk during late lactation. However, by 90 days in milk (DIM), healthy cows consistently outperformed metritic cows in terms of milk vield.

Clinical and puerperal metritis both have significant adverse effects on peak milk yield, especially during early lactation, with puerperal metritis having a more pronounced impact on production than clinical metritis (Dubuc et al., 2011). Fourichon et al. (2000) emphasized the detrimental effect of metritis on fertility, including delayed return to cyclicity, disruption of the uterine environment, and impaired embryo development (Sheldon et al., 2002; Sheldon and Dobson, 2004; Soto et al., 2003). To reduce the economic burden of metritis, it is essential for dairy farms to implement health-monitoring programs that detect and treat cows with metritis promptly, thereby minimizing its adverse productivity effects on and reproductive performance.

3. FACTORS CAUSING METRITIS

Metritis, an inflammatory condition of the uterus, is influenced by a variety of factors, many of which are linked to parturition and the early postpartum period. Factors such as parity, calving month, infectious disease presence, particularly metritis, and abnormalities during parturition significant play roles in its development. Wenz et al. (2011) found that primiparous cows and those diagnosed with contagious diseases like metritis exhibited higher rectal temperatures (RT), with peak RT occurring on the third day postpartum. These findings align with the fact that normal postpartum RT may be higher during warm summer months compared to typical values (Wenz et al., 2011).

The development of metritis is predisposed by several physical and environmental factors. These include dystocia, twin births, retained fetal placenta), membranes (retained stillbirths. hygiene deficiencies, and metabolic imbalances around parturition, such as low dry matter intake (DMI) and high prepartum serum non-esterified fatty acid (NEFA) concentrations (Ospina et al., 2010). Common pathogens associated with metritis include Arcanobacterium (Actinomyces) Trueperella pvoaenes. pyogenes, coliform bacteria, and anaerobic species such as Fusobacterium necrophorum, Prevotella spp., and Bacteroides spp. (Sheldon and Dobson, 2004; Williams et al., 2005). These bacteria necessitate antibacterial treatment to manage the condition. Historically, it was believed that the uterus was sterile before calving, and that postpartum metritis in dairy cows was caused by the introduction of Escherichia coli (E. coli) into the uterus during or after parturition. This theory suggested that E. coli facilitated secondary bacterial growth, contributing to metritis (Sheldon et al., 2009). However, more recent studies have challenged this belief, showing that both healthy and metritic cows share most bacterial genera at the time of parturition and on day six postpartum, with significant differences primarily in the abundance of certain species (Jeon et al., 2015). The uterine microbiota predominantly comprises Bacteroidetes. Fusobacteria. Proteobacteria. Tenericutes, and Firmicutes, irrespective of infection status (Santos and Bicalho, 2012; Jeon et al., 2015). Shifts in the growth of specific bacteria, such as Bacteroides, Porphyromonas, and Fusobacterium, have been linked to the onset of metritis (Jeon et al., 2015; Bicalho et al., 2017a). Notably, E. coli presence in metritic cows is low (less than 1%), and its role in metritis is only significant in the presence of virulent strains carrying the fimH factor (Bicalho et al., 2012; Jeon et al., 2015, 2016).

Primiparous cows, those experiencing abnormal calving (e.g., dystocia), and those with poor energy balance prepartum, as evidenced by elevated NEFA and low IGF-1 levels, face a higher risk of metritis (Giuliodori et al., 2013). This aligns with findings by Kaneene and Miller (1995) and Dubuc et al. (2011), who found that primiparous cows are at a higher risk due to the increased likelihood of requiring calving assistance, which increases the risk of uterine contamination and lesions (Ghavi Hossein-Zadeh and Ardalan, 2011). Studies have identified specific NEFA thresholds for predicting metritis; Ospina et al. (2010) suggested a threshold of

300 µM, while Dubuc et al. (2011) reported a threshold of 600 µM. Giuliodori et al. (2013) proposed a prepartum NEFA cut-off of 431 µM. In addition to elevated NEFA levels, other metabolic imbalances are significant predictors of metritis. For instance, Torres et al. (2020) found that cows with β -hydroxybutyrate (BHBA) concentrations \geq 0.8 mmol L^A-1 were 2.2 times more likely to develop metritis and 4.4 times more likely to develop clinical endometritis (CE). Elevated serum BHBA levels indicate poor adaptation to the demands of lactation and negative energy balance, which impair immune function (Zarrin et al., 2014). The optimal BHBA cut-off for predicting puerperal metritis was found to be 1.2 mmol LA-1. Furthermore, cows with serum creatinine concentrations $\geq 2.0 \text{ mg dL}^{-1}$ were at higher risk for metritis and CE due to prolonged tissue protein catabolism impairing immune responses (Titgemeyer and Loest, 2001). Higher serum total protein (TP) levels (≥ 5.0 mg dL^-1) were also associated with increased odds of metritis, suggesting an activated immune response post-infection (Cray et al., 2009).

Additional factors, such as body condition score (BCS), play a role in metritis risk. A BCS of \leq 3.2 should be avoided to prevent endometritis (Galvão et al., 2009). Cows experiencing physiological. significant metabolic. and immunological changes during early lactation are also more susceptible to diseases like metritis and endometritis (Bicalho et al., 2017b). High plasma glucose levels during early lactation have been associated with an increased risk of postpartum diseases. includina metritis. Furthermore, excessive loss of body energy reserves post-calving impairs immune function, making cows more vulnerable to disease, especially under high ambient temperatures (Thompson et al., 2014). Finally, cows diagnosed with metritis are at higher risk for developing clinical and cytological endometritis (Galvão et al., 2009; Martinez et al., 2012). The severity of the immune response in metritic cows is more closely correlated with disease severity than the intrauterine microbiome itself (Jeon et al., 2016). Factors such as parity, retained placenta, calving abnormalities (e.g., dystocia, twins, vulvovaginal lacerations), and elevated metabolic markers (NEFA, BHBA) have been identified as risk factors for metritis and may have predictive value for cure (Lima et al., 2014; Vergara et al., 2014; Vieira-Neto et al., 2016). The circulating concentration of haptoglobin (Hp), an acute phase protein, has also been linked to metritis

severity, suggesting its potential role as an indicator of the likelihood of cure (Huzzey et al., 2009). Thus, metritis development is influenced by a combination of cow-related factors, metabolic imbalances, and infectious agents. Early identification and management of these risk factors are crucial for preventing metritis and mitigating its consequences on dairy cattle health and productivity.

4. TEMPERATURE CHANGE MONITORING DURING METRITIS

Rectal temperature (RT) is a critical parameter in diagnosing and monitoring the progression of metritis in dairy cows. Smith et al. (1998) defined a rectal temperature exceeding 39.2°C as indicative of fever in the context of metritis. Several studies have employed different RT thresholds to assess metritis, with values ranging from 39.2°C (Smith et al., 1998) to 39.7°C (Sheldon et al., 2004), depending on the study context. Drillich et al. (2001) used a threshold of 39.5°C, while Kristula et al. (2001) recommended 39.4°C for primiparous cows and 39.7°C for multiparous cows. These variations reflect differences in study design, cow parity, and other influencing factors. A meta-analysis by Garzon et al. (2022) found that the most commonly used rectal temperature threshold for diagnosing metritis was ≥39.5°C, used in 56.8% of studies, followed by ≥39.2°C, which was used in 2.8% of the studies. This highlights the widespread consensus on the ≥39.5°C threshold for identifvina fever associated with metritis. Additionally, Garzon et al. (2022) reported variations in the methods used for evaluating vaginal discharge, with rectal palpation being the most frequently used approach (37.3%), followed by intravaginal exploration with a gloved hand (18.4%), Metricheck (9.8%), and speculum examination (5.7%). In 28.7% of the studies, diagnostic tools used for evaluating vaginal discharge were not specified.

The RT of cows diagnosed with infectious postpartum diseases, including metritis, tends to fluctuate significantly during the first ten days in milk (DIM). Burfeind et al. (2010) confirmed that repeatable and accurate RT measurements are crucial for diagnosing metritis, and noted that thermometer type and depth of insertion into the can influence individual rectum RT measurements. In their study, Wenz et al. (2011) suggested an optimal RT measurement depth of 11 cm. The study also observed that primiparous cows were more likely to develop

metritis or co-infections with other infectious diseases during the first 10 DIM compared to multiparous cows (P = 0.004; P = 0.021). Furthermore, Wenz et al. (2011) reported that cows with metritis had a peak RT of 39.6 ± 0.1°C on the day of diagnosis, which was 0.8°C higher than that of healthy parity-matched cows (P = 0.001). This elevated RT persisted for at least four days before and after the diagnosis of metritis, providing valuable insight into the temporal dynamics of RT changes in infected cows. The fever associated with metritis is primarily driven by the inflammatory response triggered by the uterine microbiome. Leukocyte activation and the release of endogenous cytokines and chemokines induce prostaglandin E synthesis, which in turn leads to fever by affecting the central nervous system (Netea et al., 2000). Both Gram-negative and Gram-positive bacteria can contribute to fever. with lipopolysaccharides (LPS) from Gramnegative bacteria being the most potent fever-inducing agents (Tavares et al., 2005). In cases characterized metritis bv an overrepresentation of **Bacteroidetes** and Fusobacteria, Gram-negative bacteria are often implicated in the fever response (Jeon et al., 2015).

Despite its utility, RT alone is not always a reliable diagnostic tool for metritis due to its overlap with other infectious diseases like mastitis. However, it remains an important parameter when used alongside other diagnostic criteria (Sheldon et al., 2006). Studies have proposed RT thresholds ranging from 39.2°C to 39.7°C for diagnosing metritis. For example, Giuliodori et al. (2013) found a receiver operating characteristic (ROC) area of 0.604 for RT at the time of metritis diagnosis, with a of 39.2°C threshold showing reasonable sensitivity and specificity. Thus, RT changes over time can provide valuable insights into disease severity and the effectiveness of treatment (Machado et al., 2020). In addition to RT, factors such as plasma haptoglobin levels, dystocia history, and milk yield can also influence temperature readings and help refine the diagnosis (Machado et al., 2020). For example, cows with a history of dystocia may exhibit higher RT due to uterine trauma, which could contribute to a more severe inflammatory response. Moreover, cows with higher milk yields might show distinct RT patterns due to increased metabolic demands post-calving, which could potentially exacerbate the risk of metritis.

Metritis is not unique to dairy cattle. In other livestock species, such as goats, metritis is often associated with retained placenta or uterine trauma and can result in infertility if not managed appropriately. In dogs and cats, metritis commonly follows prolonged labor and retained fetal tissues, with symptoms including fever, lethargy, and vulvar discharge. Mares are particularly susceptible to septic metritis, a lifethreatening condition that requires prompt intervention, especially in larger draft breeds. Similar to dairy cows, these species benefit from early detection and temperature monitoring as part of a comprehensive diagnostic approach. A promising new method for monitoring RT in dairy cows is infrared thermography (IRT), which offers a non-invasive and remote way to detect elevated RT. IRT has shown potential for early of metritis, detection enabling proactive management and improving herd health (Skliarov et al., 2023). By using RT alongside other clinical parameters, such as body condition score (BCS), leukocyte counts, and metabolic markers, dairy farmers can better manage metritis and mitigate its impact on cow health and productivity.

5. DIAGNOSTIC METHODS FOR METRITIS

Metritis remains one of the most inconsistently diagnosed and documented diseases in commercial dairy farming (Sannmann et al., 2012). Diagnostic challenges include variability in definitions, inconsistent approaches by herd personnel, and differences in the timing of disease evaluation (Espadamala et al., 2016). These inconsistencies lead can to misclassification, underdiagnosis or which introduces bias and underestimates the association between disease and performance outcomes (Johnson et al., 2014). Accurate and consistent diagnostic methods are critical for advancing our understanding of metritis' impact on productivity and improving herd health management practices.

Clinical Examination and Vaginal Discharge Evaluation: The diagnosis of metritis typically relies on clinical examination, focusing on uterine enlargement and the characteristics of vaginal discharge. Common clinical signs include a fetid, watery discharge, systemic symptoms such as fever, depression, anorexia, and dehydration, and an enlarged uterus (Benzaquen et al., 2007; Risco and Melendez Retamal, 2011). Espadamala et al. (2018) reported significant variability in diagnostic practices across dairy

farms, with discrepancies in how systemic illness and discharge characteristics were evaluated. This lack of standardization underscores the need for universally accepted diagnostic protocols to reduce the risk of underdiagnosis and ensure accurate identification.

Vaginal discharge scoring systems have been used to categorize discharge characteristics on scales such as the modified Vaginal Discharge (VD) scoring system (Oliveira et al., 2020), which ranges from 1 (normal) to 5 (fetid discharge indicative of metritis). Metricheck, a widely used tool, facilitates easy collection of vaginal discharge samples and has shown higher sensitivity but lower specificity compared to other methods (McDougall et al., 2011). Garzon et al. (2022), in their meta-analysis, highlighted the variability in descriptors for vaginal discharge color, odor, and viscosity across studies, further complicating consensus in metritis diagnosis. Terms like "fetid," "putrid," and "purulent" were commonly used but lacked consistent definitions, need for standardized emphasizing the terminology.

Rectal temperature (RT) is a key diagnostic parameter for detecting puerperal metritis (PM). Studies have used different RT thresholds for diagnosis, with ≥39.5°C being the most commonly cited cut-off (Garzon et al., 2022; Benzaguen et al., 2007). Daily RT monitoring is critical, as fever typically precedes clinical signs. For example, cows diagnosed with PM exhibit linear increases in RT starting 48 hours before diagnosis, peaking at ≥39.6°C (Benzaguen et al., 2007; Wittrock et al., 2011). Despite its utility, RT alone may not be reliable due to overlapping symptoms with other diseases, such as mastitis. Kristula et al. (2001) noted that 48% of cows with normal calving showed transient RT spikes (>39.1°C), compared to higher rates in cows with retained placenta, mastitis, PM, and dystocia. This suggests that a comprehensive health monitoring strategy combining RT measurements with other diagnostic tools is essential.

Systemic and Automated Monitoring Tools: Emerging technologies such as automated activity and rumination monitors show promise in metritis detection. Stangaferro et al. (2016) demonstrated that metritic cows exhibit altered rumination and activity patterns during the first week postpartum. These changes can be captured using automated health monitoring systems, which provide real-time data on cow behaviour and health index scores. While promising, further research is needed to validate

their efficacy in commercial farm settings. Blood biomarkers like haptoglobin (Hp), an acute-phase protein, also show potential as diagnostic tools. Elevated Hp levels at 3 and 7 days in milk (DIM) have been associated with metritis severity, poor reproductive outcomes, and reduced milk yield (Pohl et al., 2015; Barragan et al., 2018). However, Hp concentrations are also elevated in other conditions, such as dystocia and retained placenta, limiting their specificity for metritis diagnosis (Huzzey et al., 2009).

Advances in Molecular and Genetic Diagnostics: Molecular diagnostics, such as multiplex PCR, offer high sensitivity and specificity for detecting uterine pathogens, including Escherichia coli, Staphylococcus aureus, and Bacillus cereus (Wei et al., 2024). This method enables simultaneous detection of multiple pathogens, reducing diagnostic time and cost compared to conventional PCR. Additionally, Sanchez et al. (2024) identified genetic factors associated with metritis susceptibility, including genes involved in metabolism and cholesterol autophagic processes, offering potential targets for genetic selection and disease management. In addition, metabolomics offers valuable insights into the uterine environment at the time of metritis diagnosis, identifying key metabolites that are strongly correlated with the overgrowth of opportunistic pathogenic bacteria. tissue damage, and immune dysregulation, thereby providing potential targets for future interventions to mitigate metritis in dairy cows (Casaro et al., 2024).

Despite advancements, diagnostic inconsistencies persist. Garzon et al. (2022) emphasized the lack of consensus in PM criteria across studies, with 40.2% of reviewed articles providing no reference for PM definitions. Standardizing diagnostic protocols, including unified scoring systems for vaginal discharge and thresholds for RT, is essential to improve the accuracy and reliability of metritis diagnosis.

6. FUTURE PROSPECTS

The future of metritis diagnosis and management in dairy herds lies in the integration of advanced technologies, precision farming practices, and holistic approaches to herd health. Emerging diagnostic tools, such as automated activity and rumination monitors, infrared thermography, and non-invasive biomarker assays, hold promise for early and accurate detection of metritis, enabling timely interventions. Molecular techniques like multiplex PCR and genetic profiling offer opportunities to identify pathogen-specific and host susceptibility factors, paving the way for targeted therapies and genetic selection for resistance. Additionally, disease developing probiotics. vaccines, and non-antibiotic therapeutics can address antimicrobial resistance concerns while improving reproductive outcomes. Standardizing diagnostic criteria and protocols globally is essential to bridge the gaps in clinical practice and research, ensuring consistency in disease management across dairy operations. Future research should focus on validating these innovative tools and approaches in real-world settings, alongside exploring economic feasibility the of implementing such technologies to improve productivity, animal welfare, and sustainability in the dairy industry.

7. CONCLUSION

Metritis remains a multifaceted challenge in dairy herd management, significantly impacting milk production, reproductive efficiency, and the economic sustainability of dairy operations. Despite advancements in diagnostic methodologies and management strategies, the of metritis requires complexity ongoing innovation and research to address persistent gaps in understanding and control. Future efforts must prioritize the development and adoption of standardized diagnostic criteria, particularly integrating non-invasive technologies. Infrared thermography and automated health monitoring systems offer promising avenues for routine, cow-side detection of rectal temperature changes and other early indicators of metritis, enhancing the accuracy and timeliness of diagnosis while animals. minimizina stress on However. genetic unravelling the molecular and underpinnings of metritis remains a critical frontier. Identifying key pathways, such as those related to cholesterol metabolism and the PI3k-Akt pathway, will provide novel therapeutic targets and support the development of precision interventions. These advancements, combined with the refinement of management practices and exploration of innovative therapeutics like probiotics and vaccines, hold the potential to transform metritis control, improving herd health, productivity. and overall dairv industrv sustainability. Through interdisciplinary research and the integration of cutting-edge technologies, the dairy sector can overcome the challenges posed by metritis, ensuring a future of animal welfare enhanced and economic viability.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

The authors are grateful to the Director, ICAR-National Dairy Research Institute (NDRI), Karnal, Haryana, India for providing all the necessary facilities to conduct the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Barragan, A. A., Piñeiro, J. M., Schuenemann, G. M., Rajala-Schultz, P. J., Sanders, D. E., Lakritz, J., & Bas, S. (2018). Assessment of daily activity patterns and biomarkers of pain, inflammation, and stress in lactating dairy cows diagnosed with clinical metritis. *Journal of Dairy Science*, 101(9), 8248– 8258.
- Bartlett, P. C., Kirk, J. H., Wilke, M. A., Kaneene, J. B., Mather, E. C. (1986). Metritis complex in Michigan Holstein-Friesian cattle: Incidence, descriptive epidemiology, and estimated economic impact. *Preventive Veterinary Medicine, 4*, 235– 248.
- Bell, M. J., & Roberts, D. (2007). The impact of uterine infection on a dairy cow's performance. *Theriogenology, 68*, 1074–1079.
- Benzaquen, M. E., Risco, C. A., Archbald, L. F., Melendez, P., Thatcher, M. J., & Thatcher, W. W. (2007). Rectal temperature, calvingrelated factors, and the incidence of puerperal metritis in postpartum dairy cows. *Journal of Dairy Science, 90*, 2804– 2814.
- Berry, D. P., Bermingham, M. L., Good, M., & More, S. J. (2011). Genetics of animal health and disease in cattle. *Irish Veterinary Journal*, 64(1), 5. https://doi.org/10.1186/2046-0481-64-5
- Bicalho, M. L. S., Machado, V. S., Higgins, C. H., Lima, F. S., & Bicalho, R. C. (2017). Genetic and functional analysis of the

bovine uterine microbiota. Part I: Metritis versus healthy cows. *Journal of Dairy Science*, *100*(5), 3850–3862. https://doi.org/10.3168/jds.2016-12058

- Bicalho, M. L. S., Machado, V. S., Oikonomou, G., Gilbert, R. O., & Bicalho, R. C. (2012). Association between virulence factors of *Escherichia coli, Fusobacterium necrophorum*, and *Arcanobacterium pyogenes* and uterine diseases of dairy cows. Veterinary Microbiology, 157(1-2), 125–131.
- Bicalho, M. L. S., Marques, E. C., Gilbert, R. O., & Bicalho, R. C. (2017). The association of plasma glucose, BHBA, and NEFA with postpartum uterine diseases, fertility, and milk production of Holstein dairy cows. *Theriogenology*, *88*, 270–282.
- Bogado, P. O., Hostens, M., Dini, P., Vandepitte, J., Ducatelle, R., & Opsomer, G. (2016). Distribution of inflammation and association between active and chronic alterations within the endometrium of dairy cows. *Reproductive Domestic Animals*, *51*(5), 751–757.
- Bromfield, J. J., Santos, J. E., Block, J., Williams, R. S., & Sheldon, I. M. (2015). Physiology and endocrinology symposium: Uterine infection: Linking infection and innate immunity with infertility in the highproducing dairy cow. *Journal of Animal Science*, 93, 2021–2033. https://doi.org/10.2527/jas.2015-8913
- Burfeind, O., Von Keyserlingk, M. A. G., Weary, D. M., Veira, D. M., & Heuwieser, W. (2010). Repeatability of measures of rectal temperature in dairy cows. *Journal of Dairy Science*, 93(2), 624–627.
- Casaro, S., Prim, J. G., Gonzalez, T. D., Cunha, F., Bisinotto, R. S., Chebel, R. C., Santos, J. E. P., Nelson, C. D., Jeon, S. J., Bicalho, R. C., Driver, J. P., & Galvão, K. N. (2024). Integrating uterine microbiome and metabolome to advance the understanding of the uterine environment in dairy cows with metritis. *Animal Microbiome, 6*, 30.
- Cray, C., Zaias, J., & Altman, N. H. (2009). Acute phase response in animals: A review. *Comparative Medicine, 59*, 517–526.
- Cunha, F., Jeon, S. J., Daetz, R., Vieira-Neto, A., Laporta, J., Jeong, K. C., Barbet, A. F., Risco, C. A., & Galvão, K. N. (2018). Quantifying known and emerging uterine pathogens, and evaluating their association with metritis and fever in dairy cows. *Theriogenology*, *114*, 25–33.

- Daetz, R., Cunha, F., Bittar, J. H., Risco, C. A., Magalhães, F., Maeda, Y., Santos, J. E. P., Jeong, K. C., Cooke, R. F., & Galvão, K. N. (2016). Clinical response after chitosan microparticle administration and preliminary assessment of efficacy in preventing metritis in lactating dairy cows. *Journal of Dairy Science, 99*(11), 8946– 8955.
- De Oliveira, E. B., Cunha, F., Daetz, R., Figueiredo, C. C., Chebel, R. C., Santos, J. E. P., Risco, C. A., Jeong, K. C., Machado, V. S., & Galvão, K. N. (2020). Using chitosan microparticles to treat metritis in lactating dairy cows. *Journal of Dairy Science, 103*(8), 7377–7391.
- De Vries, A. (2017). Economic trade-offs between genetic improvement and longevity in dairy cattle. *Journal of Dairy Science, 100*, 4184–4192.
- Drillich, M., Beetz, O., Pfutzner, A., Sabin, M., Sabin, H. J., Kutzer, P., Nattermann, H., & Heuwieser, W. (2001). Evaluation of a systemic antibiotic treatment of toxic puerperal metritis in dairy cows. *Journal of Dairy Science, 84*, 2010–2017.
- Dubuc, J., Duffield, T. F., Leslie, K. E., Walton, J. S., & LeBlanc, S. J. (2010). Risk factors for postpartum uterine diseases in dairy cows. *Journal of Dairy Science*, *93*(12), 5764–5771.
- Dubuc, J., Duffield, T. F., Leslie, K. E., Walton, J. S., & LeBlanc, S. J. (2011). Effects of postpartum uterine diseases on milk production and culling in dairy cows. *Journal of Dairy Science*, 94, 1339–1346.
- Espadamala, A., Pallarés, P., Lago, A., & Silva-Del-Río, N. (2016). Fresh-cow handling practices and methods for identification of health disorders on 45 dairy farms in California. *Journal of Dairy Science, 99*, 9319–9333.

https://doi.org/10.3168/jds.2016-11178

- Espadamala, R., Pereira, P., Pallarés, P., Lago, A., & Silva-Del-Río, N. (2018). Metritis diagnosis and treatment practices in 45 dairy farms in California. *Journal of Dairy Science, 101*, 9608–9616.
- Fourichon, C., Seegers, H., & Malher, X. (2000). Effect of disease on reproduction in the dairy cow: A meta-analysis. *Theriogenology*, 53, 1729–1759.
- Gaddis, K. P., Cole, J. B., Clay, J. S., & Maltecca, C. (2012). Incidence validation and relationship analysis of producerrecorded health event data from on-farm computer systems in the United States.

Journal of Dairy Science, 95(9), 5422–5435.

- Galvão, K. N., Greco, L. F., Vilela, J. M., Sá Filho, M. F., & Santos, J. E. P. (2009). Effect of intrauterine infusion of ceftiofur on uterine health and fertility in dairy cows. *Journal of Dairy Science*, *92*, 1532– 1542.
- Garzon, A., Habing, G., Lima, F., Silva-Del-Rio, N., Samah, F., & Pereira, R. (2022). Defining clinical diagnosis and treatment of puerperal metritis in dairy cows: A scoping review. *Journal of Dairy Science*, 105(4), 3440–3452.
- Ghavi Hossein-Zadeh, N., & Ardalan, M. (2011). Cow-specific risk factors for retained placenta, metritis, and clinical mastitis in Holstein cows. *Veterinary Research Communications, 35*, 345–354.
- Gilbert, R. O. (2016). Management of reproductive diseases in dairy cows. *Veterinary Clinics: Food Animal Practice,* 32, 387–410. https://doi.org/10.1016/j.cvfa.2016.01.009
- Giuliodori, M. J., Magnasco, R. P., Becu-Villalobos, D., Lacau-Mengido, I. M., Risco, C. A., & de la Sota, R. L. (2013). Metritis in dairy cows: Risk factors and reproductive performance. *Journal of Dairy Science, 96*, 3621–3631.

https://doi.org/10.3168/jds.2012-6159

Gröhn, Y. T., Rajala-Schultz, P. J., Allore, H. G., DeLorenzo, M. A., Hertl, J. A., & Galligan, D. T. (2003). Optimizing replacement of dairy cows: Modeling the effects of diseases. *Preventive Veterinary Medicine*, 61(1), 27–43. https://doi.org/10.1016/S0167-

5877(03)00158-2

- Hammon, D. S., Evjen, I. M., Dhiman, T. R., Goff, J. P., & Walters, J. L. (2006). Neutrophil function and energy status in Holstein cows with uterine health disorders. *Veterinary Immunology and Immunopathology, 113*(1-2), 21–29. https://doi.org/10.1016/j.vetimm.2006.03.0 22
- Heikkilä, A. M., Nousiainen, J. I., & Pyörälä, S. (2012). Costs of clinical mastitis with special reference to premature culling. *Journal of Dairy Science*, *95*, 139–150.
- Huzzey, J. M., Duffield, T. F., LeBlanc, S. J., Veira, D. M., Weary, D. M., & von Keyserlingk, M. A. G. (2009). Short communication: Haptoglobin as an early indicator of metritis. *Journal of Dairy Science, 92*, 621–625.

- Huzzey, J. M., Veira, D. M., Weary, D. M., & von Keyserlingk, M. A. G. (2007). Prepartum behavior and dry matter intake identify dairy cows at risk for metritis. *Journal of Dairy Science*, *90*, 3220–3233.
- Jeon, S. J., Cunha, F., Ma, X., Martinez, N., Vieira-Neto, A., Rodrigo, C., Bicalho, S., Lima, S., Santos, J. E. P., Jeong, K. C., Galvão, K. N., Daetz, R., et al. (2016). Uterine microbiota and immune parameters associated with fever in dairy cows with metritis. *PLOS ONE, 11*(11), e0165740.
- Jeon, S. J., Vieira-Neto, A., Gobikrushanth, M., Daetz, R., Mingoti, R. D., Parize, A. C., et al. (2015). Uterine microbiota progression from calving until establishment of metritis in dairy cows. *Applied and Environmental Microbiology*, *81*, 6324–6332. https://doi.org/10.1128/AEM.01818-15
- Johnson, C. Y., Flanders, W. D., Strickland, M. J., Honein, M. A., Howards, P. P., Kaneene, J. B., & Miller, R. (1995). Risk factors for metritis in Michigan dairy cattle using herd- and cow-based modelling approaches. *Preventive Veterinary Medicine*, *23*(3-4), 183–200.
- Kaniyamattam, K. A., De Vries, L., Tauer, W. T., & Gröhn, Y. T. (2020). Economics of reducing antibiotic usage for clinical mastitis and metritis through genomic selection. *Journal of Dairy Science*, 103, 473–491.
- Kasimanickam, R., Duffield, T. F., Foster, R. A., Gartley, C. J., Leslie, K. E., Walton, J. S., & Johnson, W. H. (2004). Endometrial cytology and ultrasonography for the detection of subclinical endometritis in postpartum dairy cows. *Theriogenology*, 62(1-2), 9–23.
- Kristula, M. I., Smith, B. I., & Simeone, A. (2001). The use of daily postpartum rectal temperatures to select dairy cows for treatment with systemic antibiotics. *Bovine Practitioner, 35*, 117–125.
- LeBlanc, S. J. (2014). Reproductive tract inflammatory disease in postpartum dairy cows. *Animal*, *8*, 54–63. https://doi.org/10.1017/S17517311140005 24
- Liang, D., Arnold, L. M., Stowe, C. J., Harmon, R. J., & Bewley, J. M. (2017). Estimating US dairy clinical disease costs with a stochastic simulation model. *Journal of Dairy Science, 100*, 1472–1486.
- Lima, F. S., Vieira-Neto, A., Snodgrass, J. A., De Vries, A., & Santos, J. E. P. (2019).

Economic comparison of systemic antimicrobial therapies for metritis in dairy cows. *Journal of Dairy Science*, *102*(8), 7345–7358.

https://doi.org/10.3168/jds.2018-15383

- Lima, F. S., Vieira-Neto, A., Vasconcellos, G. S. F., Mingoti, R. D., Karakaya, E., Solé, E., Bisinotto, R. S., Martinez, N., Risco, C. A., Galvão, K. N., & Santos, J. E. P. (2014). Efficacy of ampicillin trihydrate or ceftiofur hydrochloride for treatment of metritis and subsequent fertility in dairy cows. *Journal* of Dairy Science, 97, 5401–5414.
- Machado, V. S., Bicalho, M. L., Rossi, R., Ribeiro, B. L., Lima, S., Santos, T., Kussler, A., Foditsch, C., Ganda, E. K., Oikonomou, G., Cheong, S. H., Gilbert, R.
 O., & Bicalho, R. C. (2014). Subcutaneous immunization with inactivated bacterial components and purified protein of *Escherichia coli, Fusobacterium necrophorum*, and *Trueperella pyogenes* prevents puerperal metritis in Holstein dairy cows. *PLOS ONE, 9*, e91734.
- Machado, V. S., Celestino, M. L., Oliveira, E. B., Lima, F. S., Ballou, M. A., & Galvão, K. N. (2020). The association of cow-related factors assessed at metritis diagnosis with metritis cure risk, reproductive performance, milk yield, and culling for untreated and ceftiofur-treated dairy cows. *Journal of Dairy Science, 103*(10), 9261– 9276.
- Macmillan, K., Colazo, M. G., & Cook, N. J. (2019). Evaluation of infrared thermography compared to rectal temperature to identify illness in early postpartum dairy cows. *Research in Veterinary Science, 125*, 315–322.
- Martinez, N., Risco, C. A., Lima, F. S., Bisinotto, R. S., Greco, L. F., Ribeiro, E. S., Maunsell, F., Galvão, K. N., & Santos, J. E. P. (2012). Evaluation of peripartal calcium status, energetic profile, and neutrophil function in dairy cows at low or high risk of developing uterine disease. *Journal of Dairy Science*, 95, 7158–7172.
- McArt, J. A., Nydam, D. V., & Overton, M. W. (2015). Hyperketonemia in early lactation dairy cattle: A deterministic estimate of component and total cost per case. *Journal* of *Dairy Science*, *98*, 2043–2054.
- McCarthy, M. M., & Overton, M. W. (2018). Short communication: Model for metritis severity predicts that disease misclassification underestimates projected milk production losses. *Journal of Dairy Science*, 101(6),

5434–5438.

https://doi.org/10.3168/jds.2017-14164

- McDougall, S., Hussein, H., Aberdein, D., Buckle, K., Roche, J., & Burke, C. (2011). Relationships between cytology, bacteriology, and vaginal discharge scores and reproductive performance in dairy cattle. *Theriogenology*, *76*(2), 229–240. https://doi.org/10.1016/j.theriogenology.20 10.12.024
- McLaughlin, C. L., Stanisiewski, E., Lucas, M. J., Cornell, C. P., Watkins, J., Bryson, L., et al. (2012). Evaluation of two doses of ceftiofur crystalline free acid sterile suspension for treatment of metritis in lactating dairy cows. *Journal of Dairy Science*, *95*(7), 4363–4371. https://doi.org/10.3168/jds.2012-5014
- Melendez, P., McHale, J., Bartolome, J., Archbald, L. F., & Donovan, G. A. (2004). Uterine involution and fertility of Holstein cows subsequent to early postpartum PGF2α treatment for acute puerperal metritis. *Journal of Dairy Science*, *87*(10), 3238–3246.
- Neave, H. W., Lomb, J., Weary, D. M., LeBlanc, S. J., Huzzey, J. M., & von Keyserlingk, M. A. G. (2018). Behavioral changes before metritis diagnosis in dairy cows. *Journal of Dairy Science*, 101(5), 4388–4399. https://doi.org/10.3168/jds.2017-13078
- Netea, M. G., Kullberg, B. J., & Van der Meer, J. W. (2000). Circulating cytokines as mediators of fever. *Clinical Infectious Diseases, 31*(Suppl 5), S178–S184. https://doi.org/10.1086/317612
- Oliveira de, E. B., Cunha, F., Daetz, R., Figueiredo, C. C., Chebel, R. C., Santos, J. E., Risco, C. A., Jeong, K. C., Machado, V. S., & Galvão, K. N. (2020). Using chitosan microparticles to treat metritis in lactating dairy cows. *Journal of Dairy Science*, *103*(8), 7377–7391. https://doi.org/10.3168/jds.2019-18028
- Olson, J. D. (1986). The metritis-pyometra complex. In *Current Therapy in Theriogenology*
- Ospina, P. A., Nydam, D. V., Stokol, T., & Overton, T. R. (2010). Evaluation of nonesterified fatty acids and βhydroxybutyrate in transition dairy cattle in the northeast United States: Critical thresholds for prediction of clinical diseases. *Journal of Dairy Science*, 93(2), 546–554.
- Overton, M. W., & Fetrow, J. (2008). Economics of postpartum uterine health. In

Proceedings of the Dairy Cattle Reproduction Council (pp. 39–44). Dairy Cattle Reproduction Council.

- Pérez-Báez, J., Silva, T. V., Risco, C. A., Chebel,
 R. C., Cunha, F., De Vries, A., Santos, J.
 E. P., Lima, F. S., Pinedo, P.,
 Schuenemann, G. M., Bicalho, R. C.,
 Gilbert, R. O., Rodríguez-Zas, S., Seabury,
 C. M., Rosa, G., Thatcher, W. W., &
 Galvão, K. N. (2021). The economic cost
 of metritis in dairy herds. *Journal of Dairy Science*, 104(3), 3158–3168.
 https://doi.org/10.3168/jds.2020-19125
- Plontzke, J., Madoz, L., De la Sota, R., Heuwieser, W., & Drillich, M. (2010). Prevalence of clinical endometritis and its impact on reproductive performance in grazing dairy cattle in Argentina. *Reproductive Domestic Animals, 46*(6), 520–526.
- Pohl, A., Burfeind, O., & Heuwieser, W. (2015). The associations between postpartum serum haptoglobin concentration and metabolic status, calving difficulties, retained fetal membranes, and metritis. *Journal of Dairy Science, 98*(7), 4544– 4551.
- Ribeiro, E. S., Lima, F. S., Greco, L. F., Bisinotto, R. S., Monteiro, A. P., Favoreto, M., et al. (2013). Prevalence of periparturient diseases and effects on fertility of seasonally calving grazing dairy cows supplemented with concentrates. *Journal* of Dairy Science, 96(9), 5682–5697.
- Risco, C., & Melendez Retamal, P. (2011). *Dairy* production medicine. Wiley-Blackwell.
- Risvanli, A., Tanyeri, B., Yildirim, G., Tatar, Y., Gedikpinar, M., Kalender, H., Safak, T., Yuksel, B., Karagulle, B., Yilmaz, O., Kilinc, M. A. (2024). Metrisor: A novel diagnostic method for metritis detection in cattle based on machine learning and sensors. *Theriogenology*, 223, 115–121.
- Sanchez, L., Campos-Chillon, F., Sargolzaei, M., Peterson, D. G., Sprayberry, K. A., McArthur, G., Anderson, P., Golden, B., Pokharel, S., Abo-Ismail, M. K. (2024).
 Molecular mechanisms associated with the development of the metritis complex in dairy cattle. *Genes*, *15*(4), 439.
- Sannmann, I., Arlt, S., & Heuwieser, W. (2012). A critical evaluation of diagnostic methods used to identify dairy cows with acute postpartum metritis in the current literature. *Journal of Dairy Research, 79*(4), 436– 444.

- Santos, T. M., & Bicalho, R. C. (2012). Diversity and succession of bacterial communities in the uterine fluid of postpartum metritic, endometritic, and healthy dairy cows. *PLoS ONE, 7*(12), e53048.
- Sheldon, I. M., & Dobson, H. (2004). Postpartum uterine health in cattle. *Animal Reproduction Science*, 82-83, 295–306.
- Sheldon, I. M., Cronin, J., Goetze, L., Donofrio, G., & Schuberth, H. J. (2009). Defining postpartum uterine disease and the mechanisms of infection and immunity in the female reproductive tract in cattle. *Biology of Reproduction, 81*(6), 1025– 1032.
- Sheldon, I. M., Lewis, G. S., LeBlanc, S., & Gilbert, R. O. (2006). Defining postpartum uterine disease in cattle. *Theriogenology*, *65*(8), 1516–1530.
- Sheldon, I. M., Noakes, D. E., Rycroft, A. N., Pfeiffer, D. U., & Dobson, H. (2002). Influence of uterine bacterial contamination after parturition on ovarian dominant follicle selection and follicle growth and function in cattle. *Reproduction*, *123*, 837–845.
- Sheldon, I. M., Rycroft, A. N., & Zhou, C. (2004). Association between postpartum pyrexia and uterine bacterial infection in dairy cattle. *Veterinary Record, 154*(9), 289– 293.
- Silva, J. C. C., Caldeira, M. O., Moraes, J. G. N., Ramos, I. S., Gull, T., Ericsson, A. C., Poock, S. E., Spencer, T. E., & Lucy, M. C. (2024). Metritis and the uterine disease microbiome are associated with long-term changes in the endometrium of dairy cows. *Biology of Reproduction*, *111*(2), 332–350. https://doi.org/10.1093/biolre/ioae067
- Skliarov, P., Fedorenko, S., Naumenko, S., Onyshchenko, O., Bilyi, D., Mylostyvyi, R., Vakulyk, V., & Kuraksina, L. (2023). Infrared thermography as a method of diagnosing reproductive pathologies in animals. *Multidisciplinary Reviews, 6*(2), 2023007.

https://doi.org/10.31893/multirev.2023007

- Smith, B. I., Donovan, G. A., Risco, C. A., Littell, R., Young, C., & Stanker, L. H. (1998). Comparison of various antibiotic treatments for cows diagnosed with toxic puerperal metritis. *Journal of Dairy Science*, *81*(7), 1555–1562.
- Soto, P., Natzke, R. P., & Hansen, P. J. (2003). Actions of tumor necrosis factor-α on oocyte maturation and embryonic development in cattle. *American Journal* of

Reproductive Immunology, 50(5), 380–388.

- Stangaferro, M. L., Wijma, R., Caixeta, L. S., Al-Abri, M. A., & Giordano, J. O. (2016). Use of rumination and activity monitoring for the identification of dairy cows with health disorders. Part III: Metritis. *Journal of Dairy Science*, *99*(9), 7422–7433.
- Stojkov, J., von Keyserlingk, M. A. G., Marchant, J. N., & Weary, D. M. (2015). Assessment of visceral pain associated with metritis in dairy cows. *Journal of Dairy Science*, *98*(8), 5352–5361. https://doi.org/10.3168/jds.2014-9296
- Tavares, E., Maldonado, R., Ojeda, M. L., & Miñano, F. J. (2005). Circulating inflammatory mediators during the start of fever in the differential diagnosis of gramnegative and gram-positive infections in leukopenic rats. *Clinical and Diagnostic Laboratory Immunology, 12*(10), 1085– 1093.
- Thompson, I. M. T., Tao, S., Monteiro, A. P. A., Jeong, K. C., & Dahl, G. E. (2014). Effect of cooling during the dry period on immune response after *Streptococcus uberis* intramammary infection challenge of dairy cows. *Journal of Dairy Science*, *97*(11), 7426–7436.

https://doi.org/10.3168/jds.2013-7621

- Titgemeyer, E. C., & Löest, C. A. (2001). Amino acid nutrition: Demand and supply in forage-fed ruminants. *Journal of Animal Science, 79*(E. Suppl), E180–E189.
- Torres, E., Mellado, M., Leyva, C., García, J. E., Véliz, F. G., & Hernández-Bustamante, J. (2020). Serum metabolites and body condition score associated with metritis, endometritis, ketosis, and mastitis in Holstein cows. *Pesquisa Agropecuária*

Brasileira, 55, e01308. https://doi.org/10.1590/S1678-3921.pab2020.v55.01308

- Vergara, C. F., Döpfer, D., Cook, N. B., Nordlund, K. V., McArt, J. A., Nydam, D. V., & Oetzel, G. R. (2014). Risk factors for postpartum problems in dairy cows: Explanatory and predictive modeling. *Journal of Dairy Science*, 97(7), 4127– 4140. https://doi.org/10.3168/jds.2012-6440
- Vieira-Neto, A., Lima, F. S., Santos, J. E. P., Mingoti, R. D., Vasconcellos, G. S., Risco, C. A., & Galvão, K. N. (2016). Vulvovaginal laceration as a risk factor for uterine disease in postpartum dairy cows. *Journal* of Dairy Science, 99(6), 4629–4637.
- Waldron, M. R., Kulick, A. E., Bell, A. W., & Overton, T. R. (2006). Acute experimental mastitis is not causal toward the development of energy-related metabolic disorders in early postpartum dairy cows. *Journal of Dairy Science*, *89*(2), 596–610.
- Wei, S., Ding, B., Wang, G., Luo, S., Zhao, H., & Dan, X. (2024). Population characteristics of pathogenic *Escherichia coli* in puerperal metritis of dairy cows in Ningxia region of China: A systemic taxa distribution of virulence factors and drug resistance genes. *Frontiers in Microbiology*, 15, 1364373.

https://doi.org/10.3389/fmicb.2024.136437 3

Wenz, J. R., Moore, D. A., & Kasimanickam, R. (2011). Factors associated with the rectal temperature of Holstein dairy cows during the first 10 days in milk. *Journal of Dairy Science*, 94(4), 1864–1872. https://doi.org/10.3168/jds.2010-3924

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/129848