



## Perception, Awareness and Treatment Experiences about Onchocerciasis in a Rural Community in Cross River State, Nigeria: Implications for Control

Regina Ejemot-Nwadiaro<sup>1\*</sup>, Emmanuel Ugobo<sup>1</sup>, Nuria Nwachuku<sup>1</sup>,  
Ukam Edadi<sup>1</sup>, Elvis Okolie<sup>1</sup>, Divine-Favour Ofili<sup>1</sup> and Glory Nja<sup>1</sup>

<sup>1</sup>Department of Public Health, Faculty of Allied Medical Sciences, University of Calabar, Calabar, Nigeria.

### Authors' contributions

This work is a research output of all authors. Authors RNE, EU, NN, UE and EO made significant input to the design of this study. Authors REN, EU, DFO and GN performed the statistical analysis. Authors REN, NN, UE and EO wrote the study protocol. Authors REN, EO, DFO, NN and GN managed the literature searches. The first draft of this manuscript was written by authors REN, EU and DFO. All authors managed the analyses and contributed to the interpretation of results of the study. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJMAH/2019/v16i230139

#### Editor(s):

(1) Dr. Nicolas Padilla-Raygoza, Professor, Department of Nursing and Obstetrics, Division of Health Sciences and Engineering, Campus Celaya-Salvatierra, University of Guanajuato, Mexico.

#### Reviewers:

(1) Hudu Okankhamame Osue, Nigerian Institute for Trypanosomiasis Research, Nigeria.  
(2) Victor Py-Daniel, Universidade de Brasília, Brasil.  
(3) O. O. Ibeh, Federal University of Technology, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50720>

Original Research Article

Received 13 June 2019  
Accepted 23 August 2019  
Published 30 August 2019

### ABSTRACT

**Background:** Onchocerciasis a disease of poverty continues to place huge health, economic and social burden on communities at risk. Understanding critical factors that impact on access to treatment, acceptance and overall control measures are pivotal to the march towards its elimination.

**Objective:** To assess Onchocerciasis perception and treatment experiences in a rural community in Cross River State, Nigeria.

**Methodology:** A cross-sectional descriptive study using mixed method was undertaken. Data were collected using pretested questionnaire and in-depth interview guide. Quantitative data were

\*Corresponding author: E-mail: [reginaejemot@yahoo.com](mailto:reginaejemot@yahoo.com);

analysed using SPSS while the in-depth interviews were audio taped, transcribed verbatim and thematic analysis done. Findings were presented in frequencies, charts, percentages, tables and quotes. Tests of significance were determined using Chi-square ( $\chi^2$ ) at significance level of 5%.

**Results:** Ignorance, myths and negative perception about the cause of Onchocerciasis pervade in 64 (31.2%) of the respondents that did not know that the bite of infected Blackfly is the cause. Some attributed the disease to curse from the gods (45.3%) and witchcraft (23.4%). This poor knowledge is associated with lower highest attained level of education ( $p = 0.01$ ). Non-availability of drugs (23.9%) and lack of knowledge on where to access ivermectin (9.8%) were the major challenges to ivermectin uptake. Inequity in treatment access was identified from the thematic analysis.

**Conclusion:** Poor knowledge of the cause of the disease, non-availability of ivermectin and misconceptions about cause of onchocerciasis had negative influence on health-seeking behaviors, discriminatory practices and treatment coverage. By integrating contextual knowledge awareness creation about Onchocerciasis into the design of control strategies will facilitate the vantage march towards achieving elimination target.

*Keywords: Onchocerciasis; access; ivermectin treatment; perception; awareness; Nigeria.*

## 1. INTRODUCTION

Onchocerciasis or river blindness is a disease of poverty that continues to place huge health, economic and social burden on communities at risk. The disease is a major problem among rural communities living in close proximity to rivers in sub-Saharan African countries. An estimated 25 million people were infected with about 1.3 million people visually impaired or blind as a result of the disease [1,2]. Nigeria is estimated to bear a significantly high burden of the disease with 32 endemic states including Cross River State [3,4]. In Cross River State, almost all the 18 local government areas (LGAs) are endemic for the disease and the prevalence was estimated to be 10% in 2012 [5], which may be gross underestimation given lack of credible population data.

Community Directed Treatment with Ivermectin (CDTI) is the major control strategy adopted in African countries by the African Program for Onchocerciasis Control (APOC). CDTI primarily involves yearly mass drug administration (MDA) of Ivermectin. Despite the successes this strategy has engendered [6,7,8], meeting target goal set for elimination of Onchocerciasis seems far-fetched [9,10,11]. However, ignorance, myths and misconceptions about Onchocerciasis have been implicated in the drug to elimination. These have equally been acknowledged to lead to negligence in prevention and control measures and causes acceptance of inappropriate treatment regimen.

It has been recognised that knowledge of history and cause of a health condition including the whole continuum of epidemiology of the disease often promotes health-seeking behaviours and encourages reduction of effects or elimination of the disease [9,11,12,13,14]. Silumbwe et al. [14] opined that often programme implementation strategies do not take into account the contextual factors that impact on overall programme success. Some of the key factors that have been suggested by many studies include; knowledge of cause and transmission of the infection, perception of disease symptoms, socioeconomic burdens of the disease, first point of call or source of treatment, factors affecting treatment regimen such as willingness to pay for treatment or otherwise, acceptance of treatment and prevention/control measures [8,11,12,13,15].

In addition, lack of knowledge of transmission of Onchocerciasis can also manifest in discriminatory and stigmatizing attitudes towards those affected [10,15]. This in turn may negatively affect the health-seeking behaviours of those affected by Onchocerciasis [8,16]. This may further limit access to ivermectin, acceptance of treatment and overall treatment coverage [13,17].

Another crucial factor could be lack of close monitoring of drug treatment and distribution by community –directed distributors (CDD) often occasioned by technical and logistic limitations in their ability to deliver interventions [1,2,5,18]. It has been equally suggested that poor compliance to treatment may not be unrelated to long treatment duration (10 – 15 years), interval between doses (one year) that can easily be

forgotten and thus missed, adverse events in ivermectin treatment often leading to rejection of treatments by communities [1,3,6]. Reinvasion caused by limited treatment coverage area has also been implicated in low CDTI programme success [15,17,19]. Perhaps this could be attributed to the inconsistent availability of ivermectin in states and government's inability to complement the efforts of APOC leading to poor distribution and follow-up in affected communities [2,5,18].

To attain community participation and design socially/locally acceptable control strategies, health program planners and implementers should be familiar with people's knowledge, attitude and practice in relation to Onchocerciasis and other cultural innuendos that impact access to treatment, coverage and other control measures [8,10]. The successful use of ivermectin at community level requires a broad public health program designed to address barriers to treatments. Understanding the

peoples' knowledge and perceptions of Onchocerciasis may stand as important promoters of effective Onchocerciasis control strategies [4,16,20]; especially in gaining the community's buy-in and confidence to participate in control programme [11,12,16,19]. There is paucity of information as few studies have been carried out to understand these issues in this environment. Therefore, this study was aimed at assessing Onchocerciasis perception, awareness and ivermectin treatment experiences among residents of a rural endemic community in Cross River State, Nigeria.

## 2. RESEARCH METHODOLOGY

### 2.1 Study Setting

The study setting was Akamkpa Local Government Area (LGA) of Cross River State in the South-South region of Nigeria (Fig. 1). It is




Fig. 1. Map of Akamkpa Local Government Area, Cross River State, Nigeria

one of the Onchocerciasis endemic foci in the State. Akamkpa LGA lies within longitude 5°25', East of the Greenwich Meridian and latitude 8°31' North of the equator. It has a projected population from the 2006 census figures to 2017 of about 203,705 using annual growth rate of 3.0%. The study area has the largest forest area in the state and a very fertile land, watered by many rivers, streams and springs; that serves as veritable breeding ground for blackflies.

## 2.2 Study Design, Sample Size and Sampling Method

This study is a cross-sectional descriptive study using a mixed method approach comprising both quantitative and qualitative data collection methods. The study population was limited to individuals residing within Akamkpa LGA of Cross River State aged 15 years and above. The sample size for this study was 205 for the quantitative data. The sample size was determined using the formula for dichotomous descriptive study [21]; employing the 10% prevalence of Onchocerciasis in Cross River State estimated by Cross River State Neglected Tropical Disease (NTD) Programme [5] at 95% confidence interval and 5% precision. Simple random sampling technique was employed to select the respondents for the quantitative aspect. A total of 25 respondents participated in the in-depth interviews comprising two from the NTD centre in Calabar, the Primary Healthcare Coordinator for Akamkpa LGA, Officers in-charge of the 10 PHCs, two active ivermectin CDDs and 10 community leaders one from each ward.

The instrument for data collection was semi-structured interviewer - administered questionnaire. It comprised of four sections. Section A elicited information on the socio-demographics of the respondent; Section B was on knowledge, perceptions and beliefs about Onchocerciasis. Both sections C and D covered Onchocerciasis treatment and factors influencing Onchocerciasis treatment respectively. In-depth Interview guide was designed to explore the experiences of participating individuals residing within Akamkpa LGA. Each interview session lasted for about 90 minutes.

## 2.3 Data Analysis

Quantitative data obtained from the study were entered, coded, cleaned and analysed using Statistical Package for the Social Sciences (SPSS version 20). Quantitative data was presented using descriptive statistics.

Categorical variables were reported as frequencies (and percentages) while normally distributed continuous variables reported as means and standard deviations. Tests of significance were determined using chi-square ( $\chi^2$ ). Each in-depth interview was tape recorded. All audiotapes from the key informants interviewed were transcribed verbatim into word documents. The transcripts and notes were analysed by themes described in the literature review as well as novel opinions expressed during the data collection process.

## 3. RESULTS

### 3.1 Socio-demographic Characteristics of Respondents

A total of 205 respondents responded to all the items in the survey questionnaire; giving a response rate of 98%. There was a slight preponderance of males; 105 (51.2%) with the respondents having a mean age of  $31.9 \pm 12.3$  years. The number of respondents were married is 103 (50.7%). Respondents with a household size between of 4- 6 were in the majority (104; 50.7%) followed distantly by respondents with 1 – 3 member household 52 (25.4%). Most of the respondents had attained secondary level of education 113 (55.1%) with those with no formal education being the least 6 (2.9%). The highest proportion of the respondents were self-employed 65 (31.7%), followed by civil servants and farmers which were equally proportioned 40 (19.5%) amongst the respondents. Most of the respondents had lived in the study area for more than 15 years 74 (36.1%). The detailed data on socio-demographic characteristics of the respondents is shown on Table 1.

### 3.2 Knowledge and Perception of Onchocerciasis

Ignorance, myths and negative perception about the cause of Onchocerciasis still persist as 64 (31.2%) of the respondents did not know that the bite of infected Blackfly is the cause (Table 2). Most attributed it to curse from the gods (29, 45.3%) and witchcraft (15, 23.4%). Having knowledge about cause of Onchocerciasis were comparatively higher in respondents who had attained more than primary level of education and reverse was the case for those that attained lower level of education (Fig. 2) was statistically significance at 0.05 critical level ( $\chi^2 = 11.32$ ;  $p = 0.01$ ). This becomes all the more significant given that majority of the respondents (55.1%)

had attained at least secondary level of education (Table 1).

Twenty four of survey respondents (11.7%) acknowledged to have suffered from Onchocerciasis. Those that were diagnosed at the health facility 16 (66.7%) and those diagnosed during mass screening exercise 8(33.3%). Having knowledge of family members suffering from the disease; only few 36 (17.6%) affirmed knowing and had one to two infected persons (51.3%) in the family (Table 2).

The Onchocerciasis prevention methods suggested by respondents (Fig. 3) were hinged on the knowledge and perception about the cause of the disease (Table 2). Among the respondents 133 (64.9%) inferred that good sanitation and personal hygiene followed by 33


(16.1%) that were of the view that wearing of protective clothing was the viable prevention strategies. Use of metizan by 8 (3.9%) and health education on prevention 5 (2.4%) were the key prevention strategies least mentioned by the respondents.

### 3.3 Access and Uptake of Ivermectin

CDDs still remain the main stay of onchocerciasis treatment (72.5%). Though health facilities (25.4%) and Patent Medicine Vendor, popularly known as “Chemist” (2.4%) were reported as the source of treatment for the rest of the respondents. A small proportion 8 (5.8%) claimed paying for the treatment (Table 2). Only very few 6(2.9%) of the respondents indicated cost of ivermectin was a challenge to its uptake (Table 3).

**Table 1. Socio demographic characteristics of respondents, Akamkpa LGA, Cross River State**


Variables	Frequency (n = 205)	Percent (%)
<b>Sex</b>		
Male	105	51.2
Female	100	48.8
<b>Family size</b>		
1 – 3	52	25.4
4 - 6	104	50.7
7- 9	35	17.1
>10	14	6.8
<b>Marital status</b>		
Single	97	47.3
Married	103	50.2
Widowed	3	1.5
Divorced	2	1.0
<b>Educational level</b>		
No formal education	6	2.9
Primary	37	18.1
Secondary	113	55.1
Tertiary	49	23.9
<b>Occupation</b>		
Civil Servant	40	19.5
Farmer	40	19.5
Self-employed	65	31.7
Student	46	22.4
Others	14	6.8
<b>Duration of stay in Akamkpa LGA</b>		
<2 years	17	8.3
2 - 5 years	28	13.7
6 – 10 years	60	29.3
11 – 15 years	26	12.7
>15 years	74	36.1
	Mean	Standard Deviation (SD)
Age (Years)	31.9	12.3



**Fig. 2. Knowledge about the cause of Onchocerciasis varied with educational level**

A significant proportion of the respondents reported having difficulties in accessing onchocerciasis treatment services (Table 3). Majority indicated that non-availability of drugs 49(23.9%) followed closely by lack of knowledge of where to get ivermectin 20 (9.8%). Other access hindering factors mentioned by

some respondents included far distance to health facility 9 (4.4%) and poor attitude of healthcare providers 9(4.4%). Possible adverse drug reaction 12(5.9%) and rejection of ivermectin 7(3.4%) were also mentioned as affecting the uptake of ivermectin.



**Fig. 3. Respondents' perception about Onchocerciasis prevention strategies**

**Table 2. Respondents' onchocerciasis knowledge and treatment profile**

<b>Variables</b>	<b>Frequency</b>	<b>Percentages</b>
<b>Knowledge of cause of Onchocerciasis</b>		
Yes	141	68.8
No	64	31.2
Total	205	100
<b>Lack of knowledge of cause of Onchocerciasis (Attributions)</b>		
Animal	9	14.1
Curse from the gods	29	45.3
Kissing	2	3.1
Witchcraft	15	23.4
Don't know	9	14.1
Total	64	100
<b>Has Onchocerciasis</b>		
Yes	24	11.70
No	181	88.29
Total	205	100
<b>How Onchocerciasis was diagnosed</b>		
Visited health facility	16	66.67
Mass screening exercise	8	33.3
Total	24	100
<b>How long with Oncho</b>		
1 – 3 months	3	12.5
4 – 6 months	2	8.33
7– 12 months	6	25.00
>12 – 36 months	5	20.83
>36 - 60 months	2	8.33
>60 months	6	25.00
Total	24	100
<b>Treatment Status (Are you on treatment?)</b>		
Yes	21	87.5
No	3	12.5
Total	24	100
<b>Source of treatment</b>		
Community Drug Distributors (CDDs)	18	85.7
Health Facility	2	9.5
Patent Medicine Store	1	4.8
Total	21	100
<b>Family member with Onchocerciasis</b>		
Yes	36	17.56
No	169	82.43
Total	205	100
<b>Number of family member with Onchocerciasis</b>		
1 – 2 persons	20	51.28
3 – 4 persons	8	22.22
5 – 6 persons	3	8.33
≥7 persons	4	11.11
Total	36	100
<b>Oncho MDA participation</b>		
Yes	138	67.3
No	67	32.9

Variables	Frequency	Percentages
Total	205	100
<b>Duration of Oncho MDA Participation</b>		
< 6 months	10	7.25
6 – 12 months	13	9.42
>12 – 36 months	51	36.96
>36 – 60 months	26	18.84
>60 months	38	27.54
Total	138	
<b>Source of Oncho MDA</b>		
Community Drug Distributors (CDDs)	100	72.5
Health Facility	35	25.4
Patent Medicine Vendor (“Chemist”)	3	2.2
Total	138	100
<b>Payment for treatment</b>		
Yes	8	5.8
No	130	94.2
Total	138	100

**Table 3. List of factors likely to affect ivermectin uptake**

Variables	*Frequency (n = 205)	
	Yes (%)	No (%)
a Drugs used for treatment not available	49 (23.9)	156 (76.1)
b Distance to the health facility is too far	9 (4.4)	196 (95.6)
c I don't know where to get the drugs	20 (9.8)	185 (90.2)
d Poor attitude of the health care providers	9 (4.4)	196 (96.6)
e Cost of drugs too high	6 (2.9)	199 (97.1)
f I don't like taking the drug	7 (3.4)	198 (96.6)
g I always forget to take my drugs as when due	6 (2.9)	199 (97.1)
h The drugs make me feel uncomfortable	12 (5.9)	193 (96.6)

\*Multiple responses; (Variables a - e speak to issues of access)

#### 4. DISCUSSION

Improving treatment access and overall coverage are critical targets that must be vigorously pursued if the set goal of elimination of onchocerciasis by year 2025 is to be achieved. However, achieving this lofty goal should be predicated on understanding critical factors that impact on treatment access, acceptance and overall onchocerciasis control measures. This study therefore sought to understand perception, awareness and treatment experiences regarding CDTI in a rural setting in Nigeria.

This study showed that about 68.8% of the respondents had knowledge about the cause of onchocerciasis to be from bite of infected black fly. This is in tandem with various studies of 69.4% in South-East Ethiopia [10] and 70% in Guatemala [15] reported knowledge levels. However, on the contrary studies by Alonso et al.

[13] in Bioko Island, Equatorial Guinea and [16] in Ogun State of Nigeria reported lower percentages of 19.3% and 9.8% respectively. This could be due to differences in educational level in the study communities.

It then follows that about 31% of the respondents in this study did not know that the bite of infected Blackfly can cause onchocerciasis. This is in spite of seemingly moderately high educational level of the respondents for most respondents (55.1%) had attained at least secondary level of education. Similarly, in a study carried out in Enugu, Nigeria, more than half of the respondents (57%) had no knowledge of the cause of onchocerciasis [12]. This thus reflects that myths and misconceptions on the cause of onchocerciasis still persist in the study area with most attributing the cause to beo curse from the gods (45.3%) and witchcraft (23.4%). This is

**Table 4. Study qualitative results**

<b>Major theme</b>	<b>Sub-themes</b>	<b>Quote</b>
Onchocerciasis is a Huge burden	Occurrence of the diseases due to the terrain, Neglected tropical diseases	“Onchocerciasis is definitely a problem; it affects the larger community in the Local Government Area”. “Yes, it a major problem as it is been called a neglected tropical disease”.
Myths and Misconceptions	Myths and Misconceptions Cause by witchcraft Curse from god Attack from enemy	“The belief in witchcraft still stands, because every small thing that happens to them, they attribute it to witchcraft”. “When people fall sick which they don’t know the possible cause they will either say it an attack from their enemy or witchcraft. “Most people in this community still belief that onchocerciasis is caused by witchcraft due to the nature of the disease”.
Discrimination and stigmatization	Negative attitude, financial incapacitation, blindness, high social burden	“You know predominantly in Akamkpa, a larger number of them are farmers, especially those in the interior, it affect them because most of them will not be able to go to Farm”. “Family that has somebody who is affected... the economy and everything in that family will not go on well, because as a father in the family you will not be able to go and fetch out what the family will eat and it will be shame and a mocking of family and stigmatization”. “it affects them because when it affects the eye, the eye is the mirror for everybody, if the eye is affected, it means even the family, community or the whole Nation is affected.” “It doesn’t actually kill but it gives indelible marks and some of them develop eye problem that they don’t know the origin”. “The economy and everything in that family will not go on well, because as a father in the family you will not be able to go and fetch out what the family will eat and it will be shame and a mocking family and stigmatization”. “The disease makes people to depend on others too much”.
Treatment of Onchocerciasis using Mectizan and Abendazole	Faith-based treatment regimen (belief, prayers)	“They are mostly treated during campaigns; we give them mectizan in combination with Abendazole mostly during campaign.” “I don’t belief the drugs work”. “Due to some peoples Religious belief, they seek the face of God or look for other alternative especially if they don’t know the possible causes”.
poor community engagement/involvement poor programme Governance and	Lack of incentives for volunteers, Poor political commitment, Religious belief,	“People who work during the first phase, during the second phase, they were not be willing saying that the money given to them is not commiserate with the job.”

<b>Major theme</b>	<b>Sub-themes</b>	<b>Quote</b>
Disillusionment	poor attitude, poor road network, Hard to reach area Language barrier, Lack of community cohesion	<p>“I stopped working to give the drugs because the families were hostile”</p> <p>“There are people who are living in very remote areas that the drugs cannot reach there, bike cannot get there, others includes language barrier and religion”.</p> <p>“Our leaders think of themselves more. They don’t care”.</p> <p>“They pay them a token at the end of their services from the donor agency... There is nothing coming from the community, or PHC”.</p> <p>“Their mentality here is quite difference, even when you take a good thing to them. They will still politicize it. Immediately they see you they will ask what have you brought for us talkless of saying how to support, they will not....”.</p>
Inequity in access	increase funding, community participation, poor Availability of Drugs Increasing awareness in hard to reach community	<p>“It’s something that Government should take control because donor at a time, they may opt out. Like in other programs that we have... if it is Government own it will be sustainable”.</p> <p>“Distribution shouldn’t be only during campaign.”</p> <p>“People should be aware; all those remote area, we should try as much as possible to reach out to them so that the people should be aware.”</p> <p>“They can step down to the community, we have to meet the opinion leaders in the community, the elders also the religious leaders especially those churches that their religion serves as a barrier.”</p> <p>“People from the Cameroon as they move in they should be able to access the drugs, So I think it should be drug that should be in the facility as they come they find it.”</p>

similar to the study carried out by [10]. Hence, observed misconceptions, myths, poor attitude and practices toward predisposing factors for onchocerciasis infection in the study area. Erroneous beliefs about onchocerciasis could lead to abandonment of personal protective measures and other preventive practices [5,8,9,10].

The pervading ignorance and poor perception on onchocerciasis is evidently reflected with most (64.9%) reported that good sanitation and personal hygiene against the small proportion that suggested use of Mectizan (3.9%) and health education on prevention (2.4%) as viable onchocerciasis prevention strategies. These x-ray the intertwined effects of lack of knowledge in reinforcing inappropriate health-seeking behaviors that invariably influence treatment distribution, acceptance and coverage [8,11,13].

Lack of knowledge and poor perception of onchocerciasis may equally not only manifest in discriminatory and stigmatizing attitudes and practices with the consequential drive for poor health-seeking behaviors that further limit access to mass drug (ivermectin) administration (MDA) [4,17], but may also affect overall efficacy of ivermectin treatment, treatment coverage and communities participation in onchocerciasis control programme [11,12,16,19]. These perceptions and ignorance were also re-echoed as major themes from the key-informants interviewed;

*“Most people in this community still believe that onchocerciasis is caused by witchcraft due to the nature of the disease” (Key informant).*

*“When people fall sick which they don’t know the possible cause they will either say it is an attack from their enemy or witchcraft” (Key informant).*

It was suggested that more than formal education may be required to bring about change that can positively influence onchocerciasis elimination target [7,11,13,16]. More importantly, the respondents are relatively young with a mean age of  $31.9 \pm 12.3$  years and ought to have access to general information often facilitated by modern technology that should be of benefit to onchocerciasis prevention and control. This therefore becomes quite pivotal in the whole scheme of onchocerciasis control, if sustained efforts at its elimination is to yield great results,

the youths as special group and this generation’s successors must be appropriately targeted with basic factual knowledge about onchocerciasis.

The few respondents that affirmed having onchocerciasis symptoms (11.1%) or family members with such symptoms (17.6%) indicated the public health burden of the disease in the study area. When the sample size ( $n = 205$ ) is matched against the population ( $N = 203,705$ ) for Akamkpa LGA as at 2017, then, the extrapolation of onchocerciasis prevalence may be far above the prevalence estimates of 10% reported in 2012 [5]. This is despite the fact that MDA of ivermectin had been on in the study area for over seven years. Thematic analysis of the qualitative aspect of this study strengthens the observation that onchocerciasis is a huge problem;

*“Onchocerciasis is definitely a problem; it affects the larger community in the Local Government Area” (Key Informant).*

*“Yes, it’s a major problem; as it is been called a neglected tropical disease” (Key Informant).*

Stigmatization, financial incapacitation and blindness were major socioeconomic variables that can have negative effects on the family, community and society from onchocerciasis. These thus strengthen the fact that onchocerciasis entrenches the vicious cycle of poverty and increases dependency. The aforementioned were listed by [4,9,12,20], that opined the association of onchocerciasis with poverty, unemployment and other social and economic consequences.

*“You know predominantly in Akamkpa, a larger number of them are farmers, especially those in the interior, it affects them because most of them will not be able to go to Farm”.*

*“Family that has somebody who is affected the economy and everything in that family will not go on well, because as a father in the family you will not be able to go and fetch out what the family will eat and it will be shame and a mocking of family and stigmatization”.*

*“It affects them because when it affects the eye, the eye is the mirror for everybody, if the eye is affected, it means even the family, community or the whole Nation is affected.”*

Respondents' indicated challenges to ivermectin uptake; non-availability of drugs (23.9%) followed by lack of knowledge of where to get the drugs (9.8%) demonstrated inequity in access to treatment. These observations are in consonance with [2,16,17,18] that inconsistent availability of ivermectin has been implicated in low Community-directed treatment with ivermectin (CDTI) programme success. It is more critical in areas with increased influx of displaced and refugee populations as was experienced in Cross River State, Nigeria.

Other factors reported by respondents include dislike for the drugs (3.4%) and fear of ivermectin-related adverse reactions (9.3%) were in agreement with [15,16] that reported fear of adverse reaction as reason for non-compliance with intake of the drugs. Adverse events in ivermectin treatment have also been acknowledged to lead to rejection of treatments by communities [1,3,6]. Thus this could limit treatment coverage and impact on possible reinvasion and perpetuate onchocerciasis endemic status of the study community.

The issue of payment for treatment (5.8%) and that of high cost of treatment (2.9%) should be source of concern in attaining elimination targets, given that CDTI are made almost entirely free-of-charge to recipients in communities at risk. Made possible by multiple source donations, coordination and collaborations [1,6,9,14,18].

## 5. CONCLUSION

Inconsistent availability of ivermectin, myths and misconceptions about cause of onchocerciasis still pervades with the dangerous consequential drive for poor health-seeking behaviours, discriminatory practices and poor treatment coverage. These findings may not be typical of the study area. The awareness of treatment and knowledge about onchocerciasis is a prerequisite for any disease control efforts. Better knowledge is shown to have a positive effect on prevention, treatment seeking and adherence to treatment, hence facilitates reductions in the socioeconomic burden of the disease. Moreover, integrating contextual knowledge about onchocerciasis in the design of control strategies could be considered as a vantage point in the march towards achieving elimination target.

## CONSENT AND ETHICAL APPROVAL

Ethical clearance to conduct this study was obtained from the Health Research Ethics Committee of Cross River State Ministry of Health. The participants were briefed on the purpose of the study and verbal consent obtained from them to enroll into the study. Participants who did not wish to be included in the research were excluded. Participants were assured of strict handling of data in confidentiality and anonymity.

## ACKNOWLEDGEMENTS

This research was supported by the African Researchers' Small Grants Program (Maiden Edition), which was funded by USAID via the Coalition for Operational Research on Neglected Tropical Diseases (COR-NTD) and administered by the African Research Network for Neglected Tropical Diseases (ARNTD). The authors wish to express their appreciation and thanks to the study communities especially the clan heads for their kind participation and local authorities particularly; to Dr Thomas Ibang, the Cross River State Coordinator NTD Programmes, Mrs E. Etene, the Primary Health Care Coordinator, Akamkpa Local Government Area (LGA), Ntufam Clement Asim and Mr Ikum Esira Ikum, the Akamkpa LGA NTD program manager and Social Mobilization Officer respectively for their continuous support throughout the study. We would also like to express our gratitude to the field team members (community volunteers) for their deep commitments to the successful completion of the project.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Centers for Disease Control and Prevention (CDC) Onchocerciasis; 2014;. Available:[https://www.cdc.gov/parasites/onchocerciasis/health\\_professionals/index.html](https://www.cdc.gov/parasites/onchocerciasis/health_professionals/index.html)
2. Weldegebreal F, Medhin G, Weldegebriel Z, Legesse, M. Knowledge, attitude and practice of community drug distributors' about onchocerciasis and community directed treatment with ivermectin in

- Quara District, North Western Ethiopia. BMC Research Notes. 2016;9(1):206.
3. Opara KN, Fagbemi BO, Atting IA, Oyene UE, Okenu DM. Status of forest Onchocerciasis in the lower Cross River Basin Nigeria: Change in clinical and parasitological indices after six years of Ivermectin intervention. Public Health. 2007;121:202–207.
  4. Umoke PC, Umoke M, Ene CU, Arua CC, Ede M. Perceived economic effects of onchocerciasis disease in Ebonyi State, Nigeria: Community health counselling implication. International Journal of Applied Engineering Research. 2018;13(21): 15136-15142.
  5. Eyo KD. Knowledge of preventive measures of onchocerciasis among adult residents of Aningeje, Akamkpa Local Government Area of Cross River State, Nigeria. Unpublished Research Project, Department of Public Health, University of Calabar, Nigeria; 2016.
  6. WHO African Programme for Onchocerciasis Control (APOC). Rapid epidemiological mapping of Onchocerciasis in Nigeria; 2005. Available: <http://www.who.int/apoc/onchoerciasis/en/>
  7. Hotez PJ. Control of Onchocerciasis: The next generation. Lancet. 2007;369(9575): 1979- 80.
  8. Dissak-Delon FN, Kamga GR, Humblet PC, Robert A, Souopgui, J, Kamgno J, Godin I. Adherence to ivermectin is more associated with perceptions of community directed treatment with ivermectin organization than with onchocerciasis beliefs. PLoS Neglected Tropical Diseases. 2017;11(8):e0005849.
  9. Charles JO, Ikpeme BM, Olaniran NS, Akpan AO, Charles AO, Ikoh MO. Biomedical paradigm and cultural perception of onchodermatitis in rural communities, Cross River State, Nigeria. African Journal of Public Health. 2007; 1(1):57–60.
  10. Weldegebreal F, Medhin G, Weldegebriel Z, Legesse M. Assessment of community's knowledge, attitude and practice about onchocerciasis and community directed treatment with Ivermectin in Quara District, north western Ethiopia. Parasites & Vectors. 2014;7(1):98.
  11. Wogu MD, Okaka CE. The knowledge, attitude and perception of onchocerciasis and ivermectin treatment by the people in Okpuje, Edo State, Nigeria. International Journal of Biomedical and Health Sciences; 2008;4(3):121-125.
  12. Ibe O, Onwujekwe O, Uzochukwu B, Ajuba M, Okonkwo P. Exploring consumer perceptions and economic burden of onchocerciasis on households in Enugu state, south-East Nigeria. PLoS Neglected Tropical Diseases. 2015;9(11): e0004231.
  13. Alonso LM, Ortiz ZH, Garcia B, Nguema R, Nguema J, Ncogo P, et al. Knowledge, attitudes and practices towards onchocerciasis among population in Bioko Island, Equatorial Guinea; Annals of Tropical Medicine and Public Health. 2017; 10(5):1228–1237. DOI: 10.4103/ATMPH.ATMPH\_726\_16
  14. Silumbwe1 A, Zulu JM, Halwindi H, Jacobs C, Zgambo J, Dambe R, Chola M, Chongwe G, Michelo C. A systematic review of factors that shape implementation of mass drug administration for lymphatic filariasis in sub-Saharan Africa; BMC Public Health. 2017;17:484. DOI: 10.1186/s12889-017-4414-5
  15. Richards Jr FO, Klein RE, de León O, Mendizábal-Cabrera R, Morales AL, Cama V, Rizzo N. A knowledge, attitudes and practices survey conducted three years after halting ivermectin mass treatment for onchocerciasis in Guatemala. PLoS Neglected Tropical Diseases. 2016;10(6): e0004777.
  16. Surakat OA, Sam-Wobo SO, Ademolu KO, Adekunle MF, Adekunle ON, Monsuru AA, Ososanya A. Assessment of community knowledge and participation in onchocerciasis programme, challenges in ivermectin drug delivery, distribution and non-compliance in Ogun State, southwest Nigeria. Infection, Disease & Health; 2018.
  17. Brieger WR, Otusanya SA, Oke GA, Oshiname FO, Adeniyi JD. Factors associated with coverage in Community-directed treatment with Ivermectin for onchocerciasis control in Oyo State, Nigeria. Tropical Medicine and International Health. 2002;7(1):11-18.
  18. Colatrella B. The Mectizan Donation Program: 20 years of successful collaboration - A retrospective. Annals of

- Tropical Medicine and Parasitology. 2008; 102(Suppl 1):7-11.
19. Braide EI, Obono MO, Basse SA. Community participation in the control of onchocerciasis in Cross River State, Nigeria. Acta Leiden. 1990;59(1-2):427-432.
20. Bawack EA. The socio-economic effects of blindness at the community and individual levels in the South-West region of Cameroon; 2018. Available: <https://www.theseus.fi/handle/10024/153526>
21. Ejemot-Nwadiaro RI. A guide to Biostatistics and health research methods. Data Pro Publishers, Calabar; 2009. ISBN: 978-051-089-3

© 2019 Ejemot-Nwadiaro et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/50720>